

Convergence Rate and Locality of Improved Overlap Fermions

W. Bietenholz

Institut für Physik
Humboldt Universität zu Berlin
Invalidenstr. 110
D-10115 Berlin, Germany

We construct new Ginsparg-Wilson fermions for QCD by inserting an approximately chiral Dirac operator — which involves ingredients of a perfect action — into the overlap formula. This accelerates the convergence of the overlap Dirac operator by a factor of 5 compared to the standard construction, which inserts the Wilson fermion as a point of departure. Taking into account the effort for treating the improved fermion, we are left with an total computational overhead of about a factor 3. This remaining factor is likely to be compensated by other virtues; here we show that the level of locality is clearly improved, so that the exponent of the correlation decay is doubled. We also show that approximate rotation invariance is drastically improved, but a careful scaling test has to be postponed.

1 Introduction

Over the recent years, substantial progress has been achieved in the long-standing problem of constructing a formulation of chiral fermions on the lattice. It turned out that there is a particularly harmless way to break the full chirality of the the lattice Dirac operator, so that the physical properties related to chirality are still represented correctly. This breaking term is sufficient to circumvent the Nielsen-Ninomiya No-Go theorem [1], which refers to full chirality, in the sense that the lattice Dirac operator D anti-commutes with γ_5 . This harmless breaking is characterized by the *Ginsparg-Wilson relation* (GWR) [2]

$$\{D, \gamma_5\}_{x,y} = 2(D\gamma_5 R D)_{x,y} , \quad (1)$$

where the kernel R has to be *local* and it must not anti-commute with γ_5 . The standard choice for this term reads

$$R_{x,y}^{st} = \frac{1}{2\mu} \delta_{x,y} , \quad \mu > 0 . \quad (2)$$

The general class of solutions to the GWR with the standard kernel R^{st} is given by the so-called *overlap formula* [3]

$$\begin{aligned} D_{ov} &= A_{ov} + \mu \\ A_{ov} &= \mu A_0 / \sqrt{A_0^\dagger A_0} \\ A_0 &= D_0 - \mu . \end{aligned} \quad (3)$$

For D_0 one may insert some sensible lattice Dirac operator. It has to be local (in the sense that the correlations decay exponentially) and free of fermion doubling, but we do not require any form of chirality for D_0 . The resulting operator A_0 is then transformed into an operator A_{ov} , so that $\frac{1}{\mu} A_{ov}$ is *unitary*. This provides the overlap Dirac operator D_{ov} , which solves the GWR. The choice of the mass parameter μ is constrained; inside its allowed interval it can be optimized with respect to certain criteria, see below.

From eq. (3) it is obvious that the spectrum of a GW fermion — with the standard kernel given in eq. (2) — is always situated on a circle in the complex plane with center and radius μ ; we denote it as the *GW circle*. This condition for the spectrum is equivalent to the GWR with kernel R^{st} .

In almost all the literature, the Wilson operator D_W was inserted as D_0 into the overlap formula, without consideration of other options. We denote

the resulting particular overlap fermion as the *Neuberger fermion*. As any Ginsparg-Wilson fermion (GW fermion), it has a chiral symmetry which is lattice modified but exact [4]. Since the corresponding transformation is local — based on the locality of R — the chiral limit is not plagued by mass renormalization.

The generalization of the overlap fermions to a whole class of solutions of the GWR — and the motivation for considering alternative operators $D_0 \neq D_W$ — were given in Ref. [5]. In fact, there are further issues of importance for a lattice fermion formulation, such as the quality of the *scaling* behavior, the level of *locality* and of approximate *rotation* invariance. With these respects, the Neuberger fermion is not quite satisfactory, even though it is local up to moderate coupling strength [6]. All these properties depend on the choice of D_0 , hence it is motivated to search for a better starting operator than D_W .

Moreover, the practical evaluation of D_{ov} is a formidable numeric task. In particular, the quenched simulation of the Neuberger fermion is about as expensive as the simulation of dynamical Wilson fermions [7], i.e. the computational overhead amounts to about two orders of magnitude. Also with this respect a better choice of D_0 can help. Our guide-line is the following observation: if we insert a GW fermion (with R^{st}) as D_0 , then it is simply reproduced by the overlap formula (3), $D_{ov} = D_0$.¹ Therefore, in practice we try to insert an approximate GW fermion as D_0 [5], so that it is altered only a little by the overlap formula,

$$D_{ov} \approx D_0 . \quad (4)$$

This small alteration corresponds to a *chiral correction* (in the sense of the GWR).²

Thus the first issue is to construct an approximate GW fermion “by hand”. At this point, we recall that also perfect [2] and classically perfect fermions [9] solve the GWR. Their construction is very tedious, but in the present context we can use a relatively simple approximation, namely a *hypercube fermion* (HF). The mass renormalization is still quite strong

¹Of course, this property holds for any fixed GW kernel R , if one uses the suitably generalized overlap formula [5].

²Similarly, one could also insert an approximately chiral D_0 in the 4d space of domain wall fermions [5, 8].

for simple HFs [10, 11], which is a problem in their direct application [12]. However, here we reach the chiral limit by inserting the HF into the overlap formula.

Of course, the (classically) perfect fermion has further virtues in addition to chirality. In particular its scaling behavior is (almost) free of lattice artifacts, i.e. dimensionless ratios of physical observables are (almost) independent of the lattice spacing. Moreover the observables are also rotational invariant for perfect fermions, and rotation symmetry is approximated very well for the truncated perfect hypercube fermions, as the pion dispersion shows [10]. Since the overlap formula modifies the HF just a little — see relation (4) — we expect the good scaling and rotation behavior to persist under the chiral correction. Then we obtain an *improved overlap fermion*.

There is yet another virtue to be expected based on relation (4): the hypercube fermion is short-ranged — its free couplings are restricted to a unit hypercube on the lattice — hence we also expect a *high level of locality* for the overlap-HF (given by D_{ov} if we insert $D_0 = D_{HF}$). Due to the modest alteration, the long range couplings can be turned on just slightly, hence their exponential decay will be fast. The Wilson fermion is also short-ranged, but it changes a lot in the transition to the Neuberger fermion, so we do not have the above reason to expect a good locality. Indeed, even for the free Neuberger fermion the decay of couplings (in the separation of fermion and anti-fermion) is a rather slow exponential [5].

All these expected virtues of the overlap-HF have been tested and impressively confirmed in the 2-flavor Schwinger model [13]. Now we carry on this program to QCD. In Section 2, we first describe the construction of a suitable HF in $d = 4$, and we illustrate its proximity to a GW fermion by evaluating typical fermionic spectra on small lattices. In Section 3 we discuss the polynomial evaluation of the inverse square root in eq. (3). In Section 4 we investigate the speed of the numeric evaluation of the overlap formula. We show that for our HF a polynomial of a low order is sufficient to approximate the sign function or the inverse square root to a high accuracy. For the Neuberger fermion the same accuracy requires a polynomial of a much higher degree. Section 5 gives results for the condition numbers and their impact on the convergence rate, now proceeding to larger lattices. Section 6 presents a comparative study of the level of locality, i.e. of the exponential decay of the maximal correlation at large distances, and Section 7 compares the violations of rotation invariance. Section 8 contains a summary and our

conclusions.

2 Approximate Ginsparg-Wilson fermions in QCD

For free or perturbatively interacting fermions, perfect actions can be constructed analytically [14]. In the case of non-perturbative interactions, this is possible only numerically and only in the classical approximation.³ The renormalization group transformation leading to the (quantum) perfect action would involve effectively a continuum functional integral. Still its existence is of conceptual interest. It implies for instance that also a topological lattice charge without lattice artifacts exists [16], and that supersymmetry can in principle be represented continuously on the lattice [17]. Similarly, the perturbatively perfect action shows that e.g. the axial anomaly is reproduced correctly [18], but only a modest improvement persists on the non-perturbative level [12, 19]. What is (in principle) accessible and promising for simulations is an approximation to the classically perfect action [20], which works well for fermions in two dimensions [21]. However, the difficult construction and application in $d = 4$ is still under investigation [22]. In that case, there is also a potential for an excellent scaling, but in order to obtain a sufficient chiral quality, it seems that the group working on it (the Bern collaboration) also depends on the concept of Refs. [5, 13] (chiral correction).

In the free case, the truncated perfect HF has still an excellent scaling behavior [10, 23], if the renormalization group parameters are optimized for locality. Hence we use the free HF as the point of departure in our attempt to construct a HF, which is an approximate GW fermion, and which is also promising with respect to scaling and approximate rotation invariance. A few elements for its gauging are then added such that the GWR is violated only modestly at the coupling strength of interest. In this Section, we are going to describe this construction step by step. A synopsis of this construction has been anticipated in Refs. [24].

The concept formulated in Refs. [5, 13] has also been adapted in Ref. [25], where some progress is reported, although a very simple (“planar”) operator D_0 was used, which is quite far from a GW fermion even in the

³For recent work on perfect actions for semi-classical effective actions, see Ref. [15].

free case. With this “planar overlap operator” some results for the finite size scaling of the chiral condensate of Ref. [26] were reproduced, and the effects of instantons in the chiral symmetry breaking were reconsidered [27]. Other very simple non-standard operators D_0 were used in Refs. [28, 29]. In contrast, a very complicated approximate Ginsparg-Wilson fermion was constructed in Ref. [30] by introducing many parameters and tuning them for a minimal GWR violation at a specific value of β . This corresponds to the first part of our program (constructing an approximate GW fermion by hand), but since the overlap formula is not used, chirality is still not exact. Moreover, there are no ingredients in favor of improving other properties. However, that work reports some gain if a specific improved gauge action is used. Different improved gauge actions were applied in Refs. [22, 31]. The use of improved gauge actions is also on our agenda, but in the present work we always use the standard plaquette gauge action. This allows us to observe unambiguously the progress due to the improved Dirac operator.

All the considerations below are based on quenched configurations on periodic lattices of sizes 4^4 , 8^4 (Sections 2, 3, 4) and 12^4 (Sections 5, 6 and 7), and beyond Section 2 we always use $\beta = 6$.

Step 1: Minimal gauging

As we mentioned above, we start from the couplings of the free perfect fermion. It is optimized for locality, and then truncated to a HF by imposing periodic boundary conditions. We write the resulting Dirac operator as

$$D(x - y) = \rho_\mu(x - y)\gamma_\mu + \lambda(x - y) , \quad (x, y \in \mathbb{Z}^4, |x_\nu - y_\nu| \leq 1) \quad (5)$$

and the couplings of the vector term ρ_μ and the scalar term λ are given in Ref. [10], Table 1. Note that the components of $x - y$ are restricted to $-1, 0, 1$, and that ρ_μ is odd in the μ -direction and even otherwise, while λ is entirely even. In the free case, this HF is an excellent approximation to a Ginsparg-Wilson fermion [5].

Our first HF for QCD, with a Dirac operator of the form

$$D(x, y, U) = \rho_\mu(x, y, U)\gamma_\mu + \lambda(x, y, U) , \quad (U : \text{compact gauge field}) \quad (6)$$

is now obtained by “minimal gauging” of the free HF: the sites, which are coupled in the HF formulation, are connected by shortest lattice path only.

The gauging is done by simply attaching the free coupling to these paths, divided into equal parts where several shortest paths exist.

This simple prescription already provides an excellent pion dispersion relation [10]. On the other hand, with such a gauging the HF suffers from a strong additive mass renormalization [10, 11], comparable to the Wilson fermion. We illustrate this at strong coupling in Fig. 1: it shows the full Dirac operator spectra for the Wilson fermion and the HF for a typical configuration at $\beta = 5$ on a 4^4 lattice.




Figure 1: *The fermionic spectrum for the minimally gauged HF and for the Wilson fermion, in a typical configuration at $\beta = 5$ on a 4^4 lattice.*

In such a rough gauge background, the overlap formula is only applicable if we have an excellent GW approximation to be inserted as D_0 . Fig. 1 shows that the standard Neuberger fermion, $D_0 = D_W$, certainly fails, and the minimally gauged HF cannot handle such a strong coupling either [32].

⁴

⁴Note that in such a situation it could be misleading to consider only the spectrum of $A_0^\dagger A_0$, as it is sometimes done in the literature. For the HF in Fig. 1, that spectrum alone would look quite satisfactory, even though the physically crucial left arc is missing.

Since we assume the kernel R to be of the form (2), the overlap formula performs a sort of projection of the eigenvalues onto the GW circle. This projection tends to be close to radial [33, 13]. Hence we need a statistically safe distinction between a left arc and a right arc of the spectrum of D_0 to start with, so that it can be mapped onto a GW circle in a reliable way. Here we focus on the regime close to the real axis. Eigenvalues have to cross the center as one changes topological sectors — since we define the topological charge by the index theorem⁵ — but they have to do so quickly as the deformation proceeds, so that the distinction is clear for the typical configurations at the coupling strength under consideration.

However, in the present case the freedom of choosing this center does still not help: wherever we choose μ , it will frequently happen that too many zeros occur (due to too many mappings to the left), so that the doubling problem is back, or that too many eigenvalues are mapped to the right-hand side, so that mass renormalization re-appears. Also further generalizations of R (beyond the restriction to one site) do not help in this situation.

A question of interest is at which coupling strength the applicability of the overlap formula with simple operators like D_{HF} sets in. Typical spectra on a 4^4 lattice suggest that for instance at $\beta = 5.4$ the coupling is still too strong, but at $\beta \gtrsim 5.6$ we are about to approach to safer grounds, see Fig. 2. The same is true for the Wilson fermion (see first Ref. in [24], Fig. 8). On larger lattices the minimal β is still likely to rise to about $\beta \gtrsim 5.7$.

Step 2: Critical link amplification

Worried about the strong mass renormalization, we first want to move our HF towards the chiral limit. We do so by amplifying each link variable by a factor $1/u$,

$$U_\mu(x) \rightarrow \frac{1}{u} U_\mu(x) , \quad u \lesssim 1 . \quad (7)$$

The idea is to compensate the (mean) link suppression due to the gauge field.

⁵This is a sensible definition of the topological charge on the lattice for any Ginsparg-Wilson fermion [4]. Up to moderate coupling strength, it is often close to the geometrical charge [34]. In general it also tends to agree with the charge identified from cooling the $SU(N)$ gauge configurations, especially at increasing N [35]. In the case of classically perfect fermions it corresponds to the classically perfect charge [9], and analogously for (quantum) perfect fermions.




Figure 2: *Typical spectra of the minimally gauged HF at $\beta = 5.4$ (on the left) and at $\beta = 5.6$ (on the right). In the latter case, the overlap formula seems to be applicable with mass parameter $\mu \approx 1.4$.*

This can be viewed as a generalization of the critical hopping parameter used for Wilson fermions, but it is also related to the spirit of “tadpole improvement” [36].

Fig. 3 illustrates that criticality is reached with $u \simeq 0.8$ at $\beta = 6$.⁶ We see that this already provides a decent approximation to a GW fermion. This is remarkable, because we approach the chiral limit in the most economic way, by staying within the framework of minimal gauging. We did not introduce additional lattice paths yet, hence Step 2 does not require any computational effort (once the critical value of u is determined).

Step 3: Fat links

We now want to improve the chiral quality of our HF further by going beyond minimal gauging. As a non-minimal element we introduce *fat links*. For a given configurations, we substitute each link variable according to the following scheme

$$link \rightarrow (1 - \alpha) link + \frac{\alpha}{6} \left[\sum staples \right], \quad \alpha \in \mathbb{R}, \quad (8)$$

⁶The absence of the arc very close to 0 is due to the *small size* of the lattice; it is *not* due to the choice of the lattice Dirac operator.




Figure 3: *HF spectra at $\beta = 6$ on a 4^4 lattice at critical link amplification ($u = 0.8$) for three configurations.*

before evaluating the Dirac operator. This is still an economic tool. The substituted link variable on the right-hand side is not mapped back onto the gauge group, in contrast to the APE blocking [38], and we do not iterate the substitution (8).

As a first observation, we note that the mass renormalization is *enhanced* for increasing α . This may seem counter-intuitive if one imagines that a positive α makes the configurations “appear smoother”. However, a more precise picture confirms this observation: the only coupling of range 0 is $\lambda(0, 0, 0, 0) = 1.853\dots$. The rest of the scalar term is negative, and in the free case we have $\sum_x \lambda(x) = 0$. The gauge field now suppresses the negative contributions, whereas $\lambda(0, 0, 0, 0)$ remains unchanged, so that a positive mass sets in (if we keep $u = 1$). This is already true for minimal gauging, but if we add fat links with $\alpha > 0$, this suppression of the negative part gets even stronger. A fraction of the negative couplings is now attached to staples instead of single links, and the staple suppression corresponds to the third power of the mean link suppression.

If we wanted to use fat links to move closer to the chiral limit, we had to

take $\alpha < 0$. Then the critical value of u rises, and for some strongly negative α (far below -1) it even arrives at 1, so that criticality could, in principle, be realized solely by means of fat links. However, in this case the rest of the spectrum is very far from a GW circle — in particular the upper and lower arc are far outside the GW circle — so we do not recommend negative values of α . Fig. 4 (on the left) shows this effect, and we also see that positive α are more adequate to make the shape of the spectrum circle-like. So what




Figure 4: *The effect due to the variation of staple coefficient α in the uncritical HF (on the left). On the right: the spectrum of the critical HF with and without fat links (everything at $L = 4$, $\beta = 6$).*

is really favorable to improve the proximity to a GW fermion is a positive α along with an adapted (i.e. decreased) critical value of u . A good choice is $\alpha = 0.3$, which requires $u = 0.76$ at $\beta = 6$. This reduces the radial fluctuation of the eigenvalues — the eigenvalues move closer together — and the upper and lower arc move closer to the GW circle. This is shown in Fig. 4 (on the right), which compares the spectra with and without fat links for the same configuration.

Step 4: Vector term suppression

The above picture for the mass renormalization ignores the rôle of the vector term $\rho_\mu(x, y, U)$. In fact, some tests confirmed that it has only a very modest influence on the location of the arc on the left-hand side (and also

on the right-hand side). This location is essentially determined by the scalar term, but the vector term is crucial for the height of the upper and lower arc, i.e. the term $\rho_\mu \gamma_\mu$ is responsible for the imaginary part of the spectrum.

This property will now be used for a further improvement, still without extra computational effort. We introduce different link amplification factors for the vector term and the scalar term, since they play a different rôle. We first keep the critical factor $1/u$ as an overall link amplification, but then we multiply a link suppression factor v only in the vector term. So now we multiply the links as follows:

$$\begin{aligned} U_\mu &\rightarrow \frac{v}{u} U_\mu && \text{in } \rho_\mu(x, y, U) \\ U_\mu &\rightarrow \frac{1}{u} U_\mu && \text{in } \lambda(x, y, U) , \end{aligned} \tag{9}$$

where $u \lesssim v \lesssim 1$. Still the fat links are useful to suppress the radial fluctuations of the eigenvalues, so we stay with $\alpha = 0.3$, $u = 0.76$, and the suitable vector term suppression amounts to $v = 0.92$. Now the upper and lower arc also follow the GW circle, and we obtain therefore a very satisfactory spectrum, see Fig. 5. It shows again the spectrum on a 4^4 lattice, but this time we also include part of the spectrum of a typical configuration on a 8^4 lattice. From there we show the 100 eigenvalues with the smallest real parts, thus we also visualize how the spectrum “continues” around zero.

A comment on the clover term

An obvious candidate for a next step beyond minimal gauging is the *clover term*. We performed a sequence of tests with it being added to the HF versions discussed above. We varied the clover coefficient and also considered both signs, but from spectra on 4^4 lattices we did not arrive at a clear further improvement of the GW approximation in this way. A positive coefficient has generally the effect to pull the spectrum closer to the real axis, as it was observed before for the Wilson fermion [37], hence the optimal parameter v rises a little. A positive clover coefficient does improve the arc around zero — which appears on the 8^4 lattice — a little bit, but it distorts the rest. As an example, we show the effect of the clover coefficient 0.15 in Fig. 6. One could argue that it is precisely the left arc which is physically crucial. However, we are going to insert our HF into the overlap formula, so we end up with an




Figure 5: *The HF on a 4^4 lattice with critical link amplification, fat links and a suppression of the vector term. We also show the “continuation” around 0 on a 8^4 lattice with the same parameters, i.e. $u = 0.76$, $\alpha = 0.3$, $v = 0.92$.*

exact GW fermion anyhow. In view of the convergence rate in the iterative evaluation of this formula, the maximal deviation of the HF spectrum from the GW circle matters, so one should not limit the attention to the left arc only.

Indeed, the systematic study of the transition to the overlap fermion on larger lattices — which will be presented in Section 5 — shows that a clover term with a small coefficient may help a little to speed up the convergence. So we are going to include it on the 12^4 lattice (used in Sections 5, 6 and 7), since it is computationally cheap anyhow.




Figure 6: *The effect of the clover term in the HF spectrum.*

3 Polynomial approximations of the overlap formula

In four dimensions, the evaluation of the overlap formula is a notorious numeric problem. It can only be done by some approximation of the non-analytic function involved. In particular, one tries to approximate it by a polynomial in the relevant interval. Here we apply Chebyshev polynomials for that purpose, as suggested in Ref. [6], and we consider two possible ways to apply them. For a systematic comparison of various approximation methods in the case of the Neuberger fermion, see Ref. [39]. We remark that from our experience, different types of suitable polynomials do not lead to a very different quality of the approximation (at fixed polynomial degree).

3.1 Approximations of the sign function

We first consider approximations of the sign function, which is done in one way or the other in most of the literature. One writes the overlap formula as

$$\begin{aligned} D_{ov} &= \mu \left[1 + \gamma_5 \frac{H}{\sqrt{H^2}} \right] = \mu [1 + \gamma_5 \epsilon(H)] \\ H &:= \gamma_5 (D_0 - \mu) = H^\dagger \end{aligned} \tag{10}$$

and approximates the *sign function*

$$\epsilon(x) = \begin{cases} 1 & x > 0 \\ -1 & x < 0 \end{cases} \tag{11}$$

by a polynomial. As an alternative to the Chebyshev polynomials that we are going to use here, also an “optimal rational approximation” has been suggested [40] and it was applied for instance in Refs. [41].

The eigenvalue distribution of H determines the interval which is relevant for the approximation. Since this Section is only meant to illustrate what the convergence rate depends on, we consider $\mu = 1$ for simplicity. (Later, when we study the convergence rate in detail, optimized mass parameters μ will be inserted.)

In Fig. 7 (on the left) we show the eigenvalue histograms for the cases of the Wilson fermion, $H_W = \gamma_5(D_W - 1)$, and of our preferred HF on small lattices ($\alpha = 0.3$, $u = 0.76$, $v = 0.92$), still at $\beta = 6$, on a 4^4 lattice. We see that the spectrum of H_{HF} is already sharply peaked at ± 1 , whereas for H_W the distribution is very broad.

In order to make the polynomial approximation directly applicable for all eigenvalues, we first have to re-scale the spectra so that they are entirely confined to the interval $[-1, 1]$, see for instance Ref. [42]. The outcome of the minimal re-scaling (division by the largest absolute value of an eigenvalue) is shown in Fig. 7 (on the right). We see that the spectrum of H_{HF} is not affected too much: the peaks are moved to about ± 0.7 , but there is still a large gap around zero. This gap is important, because any polynomial approximation of the sign function is plagued by its worst errors near the discontinuity at zero (remember for instance the notorious “Gibbs phenomenon” of the Fourier expansion). On the other hand, for H_W there is (after re-scaling) a considerable eigenvalue density around zero. This shows that it requires much more effort to transform the Wilson fermion into an overlap fermion.




Figure 7: *On the left: the eigenvalue histograms for typical spectra of H_{HF} (grey) and H_W (black) at $\beta = 6$. On the right: the same after re-scaling so that all eigenvalues have absolute values ≤ 1 .*

To demonstrate this prediction with an example, we use a linear combination of Chebyshev polynomials with maximal degree 21 to approximate the sign function in the overlap formula (10). We consider again a typical configuration at $\beta = 6$ on a 4^4 lattice. Fig. 8 shows the resulting spectra if we start from the Wilson fermion resp. the HF, and we see that the latter is clearly superior.

3.2 Approximation of the inverse square root

We now come to a second way to approximate the overlap formula in terms of polynomials. The idea is not to consider the sign function any more, but to approximate directly the inverse square root in eq. (3). This was first applied in Ref. [6]. If we start from an approximate GW operator D_0 , then the square root is close to 1 (or in general close to μ), and the direct expansion around this constant looks attractive. We have tested this method in the Schwinger model and we found a very fast convergence if we start from the 2d HF, but it slows down if we start from the Wilson fermion [13]. To get started with the polynomial approximation, one re-scales the operator A_0 so that the spectrum of $A_0^\dagger A_0$ is all contained in an interval $[\delta, 1]$ ($0 < \delta \ll 1$). The virtue of the function to be expanded, $f(x) = 1/\sqrt{x}$, is its continuity in this interval. However, if δ becomes very small we approach a singularity, which is a problem similar to the discontinuity of the sign function. Hence the question here is how small δ is going to be. Actually the question is the same as in Subsection 4.1 (and also the re-scaling is the same), because




Figure 8: *The spectra of approximate overlap fermions on a 4^4 lattice, where the sign function is approximated by a polynomial of degree 21. We use a configuration typical at $\beta = 6$ and start from $D_0 = D_W$ (stars) resp. from $D_0 = D_{HF}$ (crosses).*

$A_0^\dagger A_0 = H^2$. So we can refer again to Fig. 7 where one just has to square the eigenvalues.

Also the comparison of spectra looks very similar to Fig. 8, so we do not repeat it but turn to a systematic study of the convergence to an overlap fermion.

4 Convergence rate

We now look explicitly at the convergence of the overlap formula approximated by polynomials.⁷ As a measure for the deviation from the final overlap fermion, we measure the maximal deviation of any eigenvalue in the spectrum of $\frac{1}{\mu}D$ from the unit circle with center 1, $\{z \mid |z - 1| = 1\}$. For an exact GW fermion — with respect to the standard kernel (2) — the spectrum

⁷The results of this and the following Sections have been summarized before in Ref. [43].

is situated exactly on this circle, as we pointed out in Section 1.

Fig. 9 shows this maximal deviation for a typical configuration on a 4^4 lattice, as a function of the polynomial degree. We consider both, the expansion of the sign function as well as the expansion of the inverse square root. The converge rate is clearly *exponential* in the degree of the polynomial. We see further that the HF converges much faster than the Wilson fermion. For the comparison between the two polynomial expansions one has to take into account that the square root expansion refers to H^2 , whereas the sign function is expanded in H . Hence the former actually picks up a factor of 2 in the comparison of the polynomial degree, which makes the sign function look favorable, especially for the HF.




Figure 9: *The maximal radial deviation from the normalized Ginsparg-Wilson circle, evaluated from of the full spectrum on a 4^4 lattice. We compare the HF at $\mu = 1$ with the Wilson fermion at $\mu = 1.6$, in both cases expanding the inverse square root as well as the sign function.*

Since the volume considered in Fig. 9 is quite small, we now proceed to a 8^4 lattice. Here we cannot evaluate the full spectrum any more, but with the Arnoldi algorithm we can identify a selected set of eigenvalues. We thus evaluated the 100 eigenvalues with the least real part (as shown in Fig. 5

before), because they are physically most relevant, and we measured for this subset again the maximal radial deviation. In Fig. 10 we plot this maximal deviation, and also the mean deviation, again as a function of the polynomial degree. For the Wilson fermion we consider two options: we take either the free hopping parameter and optimize the mass parameter to $\mu = 1.6$, or we take $\mu = 1$, as in the case of the HF, and insert the critical Wilson fermion. The goal is to make sure that we compare the HF really to the best application of the Wilson fermion. However, as Fig. 10 shows, the difference between the different ways to use the Wilson fermion is tiny.




Figure 10: *The maximal radial deviation and the mean radial deviation from the normalized Ginsparg-Wilson circle, evaluated from of the 100 energy eigenvalues with the lowest real parts (physical branch) on a lattice volume 8^4 . We compare the inverse square root expansion for the HF and for the Wilson fermion. In the latter case we also compare the case of the free hopping parameter and a suitable mass parameter of $\mu = 1.6$ with the critical Wilson fermion at $\mu = 1$. They behave very similarly, and the HF converges much faster.*

Due to the arbitrary truncation at just 100 eigenvalues the exponential behavior is not as clean as in the case of the full spectrum. However, we

see that the behavior is very similar, and the improvement of the HF clearly persists in the same magnitude.

To make these observations more quantitative, we discuss as an example the sign expansion on the 4^4 lattice, which has a very precise exponential behavior. For some configuration typical at $\beta = 6$ we measured for the Wilson fermion the maximal deviation $d_W^{max}(n) = \exp(-0.134n)$, where n is the degree of the Chebyshev polynomial, and for the mean deviation we found $d_W^{mean}(n) = 0.13 \exp(-0.134n)$, hence the exponential factor is the same. For the same configuration, the following HF deviations were obtained: $d_{HF}^{max}(n) = \exp(-0.737n)$, and $d_{HF}^{mean}(n) = 0.1 \exp(-0.737n)$.

We mention two ways how to arrive at conclusions from these numbers.

- First, we could fix some degree n which we consider affordable in a simulation. The precision of the GW approximation compares as

$$\frac{d_W^{max}(n)}{d_{HF}^{max}(n)} \cong e^{0.6n} . \quad (12)$$

For realistic degrees like $n = 20 \dots 100$ this ratio of the accuracies takes a very considerable magnitude.

- On the other hand, we could fix a certain accuracy d^{max} which we consider necessary to trust the chiral quality of the approximated GW fermion. Then the polynomial degrees, which are required to provide this precision, compare as

$$\frac{n_W}{n_{HF}} \cong 5.5 . \quad (13)$$

This factor may be regarded as the effective gain of the HF due to the faster convergence, since the computational effort is essentially proportional to n .

The fluctuation of these ratios over different configurations are modest, even though d^{max} may vary significantly. For a systematic statistical study we now move on to larger lattices.

5 Condition numbers

After the explicit convergence study of the last Section, we now turn our attention to the condition numbers of the operators $A_0^\dagger A_0$, which are crucial for the convergence rate. This allows us to proceed to larger lattices of size 12^4 , still at $\beta = 6$. We first show a history for the condition numbers for the HF and the Wilson fermion in Fig. 11 (on top). Here we adapted the parameters so that they are optimal on the larger lattice. For the HF the new set of optimal parameters reads

$$u = 0.75, \quad v = 0.99, \quad \alpha = 0.87, \quad c = 0.01, \quad \mu = 1.3. \quad (14)$$

For the Wilson fermion (with the free hopping parameter), $\mu = 1.64$ turned out to be optimal with respect to the condition number, though the dependence on μ is weak — as it is also the case for locality, see Section 6. Hence the condition numbers at $\mu = 1.4$ — which is optimal with respect to locality [6] — differ only little from the result in Fig. 11.

We see that the HF condition number is improved typically by one to two orders of magnitude. Since it is defined by the ratio of the upper bound divided by the lower bound, it is instructive to consider these bounds separately. Their histories are shown in Fig. 11 (below). They reveal that the improvement of the condition number is almost entirely an effect due to the *upper* bound.

In Fig. 12 we illustrate explicitly how the condition number translates into an accelerated convergence. As a (somewhat arbitrary) measure for the speed of convergence, we consider Chebyshev polynomials at some moderate degree, $n = 60$, and measure the deviation of the function $f_{\max}(r = 24)$ from the exact result (obtained from huge values of n). (The quantity $f_{\max}(r = 24)$ represents the maximal correlation over the largest distance on our lattice; an explicit definition will be given in Section 6.) A polynomial of degree 60 may be affordable in simulations, and we see that it typically approximates $f_{\max}(r = 24)$ already to a high accuracy for the HF, but not for the Wilson fermion. For the latter we see that $\mu = 1.64$ is better for the condition number than the locality optimal mass parameter of $\mu = 1.4$.

However, it is important to note that most practical applications of overlap fermions are performed such that the lowest few modes are projected out and treated separately. Then the above polynomial evaluation concerns the




Figure 11: *On top: a history of the condition numbers of $A_0^\dagger A_0$ at $\beta = 6$ on a 12^4 lattice. Below: the corresponding history of the upper and lower bound of the spectra. It shows that the improvement of the condition number of the HF is essentially due to the upper bound.*

rest of the spectrum. This method helps a lot, because often very few modes are responsible for a slow convergence. However, their separate treatment is also tedious, because it requires a very accurate determination of the corresponding eigenfunctions. Hence the number of modes to be projected out is usually not more than about 15; beyond that the spectrum becomes quite dense, so projecting out further eigenvalues raises the remaining lower bound only very little.

To do justice to this situation, we introduce “higher condition numbers”




Figure 12: *The precision of the overlap formula approximated by Chebyshev polynomials of the moderate degree of 60. As a measure for the precision, we show the accuracy of the function $f_{\max}(r = 24)$ (c.f. Section 6) for a set of configurations at $L = 12$, $\beta = 6$. We plot this accuracy against the condition number, in order to illustrate their monotonous relation, and the progress of the HF over the Wilson fermion.*

c_k , which are defined by the ratio

$$c_k = \frac{\text{largest eigenvalue}}{k^{\text{th}} \text{ eigenvalue from below}} . \quad (15)$$

If one projects out 15 eigenvalues, for example, then c_{16} is relevant for the convergence of the polynomial evaluation. In Fig. 13 we show the histories for the higher conditions numbers $c_2 \dots c_{20}$, for the HF and the Wilson fermion. This plot confirms that around $k = 15$ the eigenvalue density is large already, and beyond they become even more dense. Hence it is hardly motivated to increase k much further. We also see that these histories are much smoother compared to c_1 (which was shown in Fig. 11), so here we are able to take sensible mean values. The results are shown in Fig. 14, and we see that in the relevant regime the improvement factor for the HF amounts to about 25. The convergence rate behaves like the square root of the relevant condition




Figure 13: *The history of the “higher condition numbers” $c_2 \dots c_{20}$ (defined in eq. (15)) for the operator $A_0^\dagger A_0$, built from the HF (at $\mu = 1.3$) and from the Wilson fermion (at $\mu = 1.64$).*

number.⁸ Hence we gain a factor of ≈ 5 . This is amazingly consistent with the explicit result obtained on the small lattices (but from the full spectrum) in Section 4.

6 Locality

Since there is occasionally some confusion about the term “locality” of a lattice action, we first clarify that we refer to the definition that all correlators decay at least exponentially in the distance. This is the property which is crucial for providing a safe continuum limit (since the decay width is fixed in lattice units).

It was conjectured [5] — and later proved [44] — that Ginsparg-Wilson fermions cannot be “ultralocal”, not even in the free case. This means that their couplings cannot drop to zero beyond a finite number of lattice spacings.

⁸Note that the relevant condition number corresponds to $1/\delta^2$ in the notation of Section 3.




Figure 14: *The expectation values for the higher condition numbers c_k ($k = 2 \dots 20$) (defined in eq. (15)) for the operator $A_0^\dagger A_0$, built from the HF and from the Wilson fermion. In the most popular regime (around $k = 15$) the HF gains a factor of ≈ 25 over the Wilson fermion. For the latter we also confirm that $\mu = 1.64$ is a little better for the condition number than the locality optimal parameter $\mu = 1.4$.*

However, locality in the above sense was shown for the free perfect fermion [14] and for the Neuberger fermion [4] to hold. In Ref. [5] it was shown that the truncated perfect free HF leads to an overlap-HF, which is more local than the Neuberger fermion (it has a faster exponential decay).

In the presence of gauge interactions, locality can be proved for smooth configurations, either by assuming a small upper limit for the deviation of any plaquette variable from 1 [6, 45], or by assuming that the eigenvalues of $A_0^\dagger A_0$ do not cluster densely in the vicinity of zero [6]. For realistic configurations, the exponential decay was observed statistically for the Neuberger fermion at $\beta = 6$ [6]. This property may collapse at much stronger coupling, but at some point the overlap formula is not applicable any more also for other reasons, as we discussed in Section 2. The statistical demonstration was done by showing that the “maximal correlation” between any two lattice sites, separated by a taxi driver distance r , decays exponentially in r . More

precisely, the expectation value of the function

$$f_{\max}(r) := \max_{x,y} \left\{ \|\psi(y)\| \mid \sum_{\mu=1}^4 |x_{\mu} - y_{\mu}| = r \right\} \quad (16)$$

has to decay exponentially, if a unit source is located at the (arbitrary) site x . We probed 6 sites x for each configuration, then we went through all sites y to determine the maxima. The 6 options for x were sufficient to stabilize the function $f_{\max}(r)$. (Remember that this function was mentioned before in Section 5.)

As we explained in the introduction, the property (4) suggests that the higher degree of locality of the overlap-HF may also persist in the presence of gauge interactions. In fact, in the Schwinger model a comparison of the function $\langle f_{\max}(r) \rangle$ confirmed this conjecture [13]. Here we extend this comparison to QCD on a periodic 12^4 lattice, which is the size that was also used in Ref. [6]. The use of the taxi driver metrics allows us to proceed to maximal distance 24, and the exponential decay is clearly visible.⁹ Our result is shown in Fig. 15. For the Neuberger fermion it agrees well with the result of Ref. [6] (which was also obtained at $\beta = 6$), although we used a different mass parameter. We inserted $\mu = 1.64$ which was optimal for the condition number of the Neuberger fermion (c.f. Section 5), whereas Ref. [6] used $\mu = 1.4$, which is slightly better for the locality — because it was optimized with this respect — but the difference is really small. Also for the overlap-HF we used the parameter which is optimal for the condition number, in that case $\mu = 1.3$.

We see that the overlap-HF is by far more local. To be explicit, the asymptotic decay of $\langle f_{\max}(r) \rangle$ behaves like $0.017 \exp(-0.45r)$ for the Wilson fermion, and like $0.017 \exp(-0.93r)$ for the HF. This observation suggests that the range of applicability of the overlap formula extends up to stronger coupling for the overlap-HF.

Interestingly, the very clean exponential decay sets in only in the presence of interactions. For free fermions, the decay is faster, of course, but the exponential behavior is not as exact as in Fig. 15.

⁹Of course, the decay is also exponential with respect to the Euclidean distance. There the decay is less smooth, but the asymptotic behavior can be extracted as $0.04 \exp(-|x-y|)$ for the Neuberger fermion, and as $0.7 \exp(-2 \cdot |x-y|)$ for the overlap HF.




Figure 15: *Comparison of the degree of locality for the Neuberger fermion and for the overlap HF. We see that the latter is clearly more local; the exponent differs by more than a factor of 2.*

On the technical side we mention that the convergence of $f_{\max}(r)$ at long distances requires a very precise evaluation of the inverse square root. This motivated the use of $f_{\max}(r = 24)$ as a measure for the precision of the approximations to the overlap formula, see Section 5.

7 Approximate rotation invariance

As a last comparison between the Neuberger fermion and the overlap HF, we consider the extent of the violations of rotation invariance. To quantify this property, we introduce the function $f_{\max}(|x - y|)$, which corresponds to definition (16) in Euclidean metrics, as well as the corresponding function $f_{\min}(|x - y|)$,

$$f_{\max}(|x - y|) \quad := \quad \max_{x, y} \left\{ \|\psi(y)\| \left[\sum_{\mu=1}^4 (x_{\mu} - y_{\mu})^2 \right]^{1/2} = |x - y| \right\} ,$$

$$f_{\min}(|x - y|) := \min_{x, y} \left\{ \|\psi(y)\| \sqrt{\sum_{\mu=1}^4 (x_{\mu} - y_{\mu})^2} = |x - y| \right\}. \quad (17)$$

We put a unit source on one site x and probe all other sites y for a fixed Euclidean distance. For each distance $|x - y|$ that occurs, we determine the difference

$$\langle f_{\max}(|x - y|) - f_{\min}(|x - y|) \rangle$$

which represents a measure for the violation of rotation invariance. Fig. 16 shows that this difference decays exponentially as a function of the Euclidean distance. (Again we present data from a periodic 12^4 lattice at $\beta = 6$.) For the Neuberger fermion (at $\mu = 1.64$), the asymptotic decay amounts to $0.065 \exp(-0.11 \cdot |x - y|)$. The corresponding asymptotic decay for the overlap HF (at $\mu = 1.3$) behaves as $\exp(-2.10 \cdot |x - y|)$.




Figure 16: *Comparison of the violations of the rotation invariance for the Neuberger fermion and for the overlap-HF. We see that the latter approximates rotation invariance much better. In both cases, the violations of rotation invariance decay exponentially with the distance, but for the overlap HF the exponent of the decay is almost doubled.*

8 Conclusions

We have constructed a HF for QCD, which approximates the GWR at $\beta = 6$. It involves the couplings of the truncated perfect free fermion, along with four more parameters which amplify certain links and add fat links as well as a clover term. This HF is designed to be inserted into the overlap formula, which renders its chirality exact. Since we start off in the right vicinity, the convergence under polynomial evaluations of the overlap formula is accelerated compared to the standard Neuberger fermion, which uses the Wilson fermion as point of departure. We observed consistently over lattice volumes 4^4 up to 12^4 an improved convergence rate the overlap-HF, which allows us to obtain the same chiral accuracy as the Neuberger fermion if the polynomial degree is reduced by a factor $\gtrsim 5$. This implies a correspondingly reduced computational effort.

Our experience with the numeric treatment of the HF is based on a simple implementation [11], which first computes (hierarchically) a look-up table for all the “hyperlink variables” across (spatial) diagonals inside the unit hypercube. This increases the number of $SU(3)$ matrices per site to be stored by a factor of 10 compared to the clover Wilson fermion. We further estimate that the computational overhead amounts to a factor ≈ 15 . Hence after the gain in the convergence rate we are left with an computational overhead of a factor of ≈ 3 .

However, we should emphasize that the algorithms for the overlap-HF are not optimized carefully yet, so there may still be space for a further gain due to a better implementation. Moreover, we expect a number of further virtues of the overlap-HF compared to the Neuberger fermion, which could well compensate this remaining overhead. We have shown that locality is clearly improved — the exponent of the correlation decay (in the distance between $\bar{\psi}$ and ψ) is doubled — which makes the overlap formula applicable up to stronger coupling. We have also shown that the overlap-HF approximates the rotation invariance very well, in contrast to the Neuberger fermion.

However, the really crucial question is if our expectation for an improved scaling can also be confirmed. If the overlap-HF allows for the use of a somewhat larger physical lattice spacing, then the remaining overhead could be more than compensated. This hope is based in particular on the elements of an truncated perfect actions, which are incorporated in our HF construction. Indeed, a strongly improved scaling was observed for free fermions and in the

2-flavor Schwinger model $d = 2$ [13]. We also run some tests for the meson dispersions in QCD, but it turned out that the dispersions are quite noisy, in particular for the overlap-HF. Hence we postpone the delicate question of scaling for further investigation. Interestingly, the phenomenon of more noise is also known from $O(a)$ improved Wilson fermions with a clover term. As a general property, GW fermions are also $O(a)$ improved. Hence it is conceivable that the overlap HF has an even better scaling behavior than the original HF, since the overlap formula removes the $O(a)$ artifacts.

As long as GW fermions can only be applied in the quenched approximation, the connection with chiral perturbation theory [46] seems most attractive. However, the ultimate goal is the use of dynamical Ginsparg-Wilson fermions, which appears as a tremendous task. Hence it is worthwhile studying the optimal access carefully. We hope that the current work contributes to this optimization.

Acknowledgment *I am very much indebted to Ivan Hip, who made important contributions to this work. I also thank him, as well as David Adams, Norbert Eicker, Philippe de Forcrand, Leonardo Giusti, Karl Jansen, Waseem Kamleh, Thomas Lippert, Martin Lüscher, Klaus Schilling, Rainer Sommer and Urs Wenger for useful discussions. The computations for this work were performed on the NICSE machines at the Forschungszentrum Jülich.*

References

- [1] H.B. Nielsen and M. Ninomiya, Phys. Lett. B105 (1981) 219; Nucl. Phys. B185 (1981) 20.
- [2] P.H. Ginsparg and K.G. Wilson, Phys. Rev. D25 (1982) 2649.
- [3] H. Neuberger, Phys. Lett. B417 (1998) 141, B427 (1998) 353.
- [4] M. Lüscher, Phys. Lett. B428 (1998) 342.
- [5] W. Bietenholz, Eur. Phys. J. C6 (1999) 537.
- [6] P. Hernández, K. Jansen and M. Lüscher, Nucl. Phys. B552 (1999) 363.

- [7] P. Hernández, K. Jansen and L. Lellouch, *in* “Numerical challenges in lattice quantum chromodynamics”, A. Frommer, Th. Lippert, B. Medeke and K. Schilling (eds), Springer (2000) p. 29 (hep-lat/0001008).
- [8] Y. Shamir, Phys. Rev. D62 (2000) 054513.
- [9] P. Hasenfratz, V. Laliena and F. Niedermayer, Phys. Lett. B427 (1998) 125. P. Hasenfratz, Nucl. Phys. B525 (1998) 401.
- [10] W. Bietenholz, R. Brower, S. Chandrasekharan and U.-J. Wiese, Nucl. Phys. B [Proc. Suppl.] 53 (1997) 921.
- [11] W. Bietenholz, N. Eicker, A. Frommer, Th. Lippert, B. Medeke, K. Schilling and G. Weuffen, Comput. Phys. Commun. 119 (1999) 1.
- [12] K. Orginos *et al.*, Nucl. Phys. B [Proc. Suppl.] 63 (1998) 904.
- [13] W. Bietenholz and I. Hip, Nucl. Phys. B570 (2000) 423; Nucl. Phys. [Proc. Suppl.] 83-84 (2000) 600.
- [14] W. Bietenholz and U.-J. Wiese, Nucl. Phys. B464 (1996) 319.
- [15] S. Kato, N. Nakamura, T. Suzuki and S. Kitahara, Nucl. Phys. B520 (1998) 323. S. Fujimoto, S. Kato and T. Suzuki, Phys. Lett. B476 (2000) 437. M.N. Chernodub, S. Fujimoto, S. Kato, M. Murata, M.I. Polikarpov and T. Suzuki, Phys. Rev. D62 (2000) 094506. M.N. Chernodub, K. Ishiguro, T. Suzuki, hep-lat/0204003.
- [16] W. Bietenholz, R. Brower, S. Chandrasekharan and U.-J. Wiese, Phys. Lett. B407 (1997) 283.
- [17] W. Bietenholz, Mod. Phys. Lett. A14 (1999) 51.
- [18] W. Bietenholz and U.-J. Wiese, Phys. Lett. B378 (1996) 222; Nucl. Phys. B [Proc. Suppl.] 47 (1996) 575.
- [19] W. Bietenholz and T. Struckmann, Int. J. Mod. Phys. C10 (1999) 531.
- [20] P. Hasenfratz and F. Niedermayer, Nucl. Phys. B414 (1994) 785. F. Niedermayer, P. Rüfenacht, and U. Wenger, Nucl. Phys. B597 (2001) 413. P. Rüfenacht and U. Wenger, Nucl. Phys. B616 (2001) 163.

- [21] C.B. Lang and T.K. Pany, Nucl. Phys. B513 (1998) 645.
F. Farchioni and V. Laliena, Phys. Rev. D58 (1998) 054501.
- [22] P. Hasenfratz, S. Hauswirth, K. Holland, T. Jörg, F. Niedermayer and U. Wenger, Nucl. Phys. B [Proc. Suppl.] 94 (2001) 627. P. Hasenfratz, S. Hauswirth, K. Holland, T. Jörg and F. Niedermayer, Nucl. Phys. B [Proc. Suppl.] 106 (2002) 751, 799; hep-lat/0205010. S. Hauswirth, Ph.D. thesis (hep-lat/0204015).
- [23] W. Bietenholz and U.-J. Wiese, Phys. Lett. B426 (1998) 114. W. Bietenholz, Nucl. Phys. A642 (1998) 275c.
- [24] W. Bietenholz, in Proceedings of the “International Workshop on Non-Perturbative Methods and Lattice QCD”, Guangzhou (China), X.-Q. Luo and E.B. Gregory (eds), World Scientific (Singapore, 2001) p. 3 (hep-lat/0007017). W. Bietenholz, N. Eicker, I. Hip and K. Schilling, Nucl. Phys. B [Proc. Suppl.] 94 (2001) 603.
- [25] T. DeGrand, Phys. Rev. D63 (2001) 034503.
- [26] P. Hernández, K. Jansen and L. Lellouch, Phys. Lett. B469 (1999) 198.
- [27] T. DeGrand and A. Hasenfratz, Phys. Rev. D64 (2001) 034512.
- [28] A. Boriçi, Phys. Lett. B453 (1999) 46.
- [29] W. Kamleh, D. Adams, D. Leinweber and A. Williams, hep-lat/0112041.
- [30] C. Gattringer, I. Hip and C.B. Lang, Nucl. Phys. B597 (2001) 451.
For a corresponding work in $d = 2$, see C. Gattringer and I. Hip, Phys. Lett. B480 (2000) 112.
- [31] S.J. Dong, F.X. Lee, K.F. Liu and J.B. Zhang, Phys. Rev. Lett. 85 (2000) 5051; Nucl. Phys. B [Proc. Suppl.] 94 (2001) 752. F.D.R. Bonnet, P.O. Bowman, D.B. Leinweber, A.G. Williams and J.B. Zhang, hep-lat/0202003.
- [32] W. Bietenholz, *in* “Lattice Fermions and the Structure of the Vacuum”, V. Mitrjushkin and G. Schierholz (eds.), Kluwer Academic C 553 (2000) p. 77 (hep-lat/0001001).

- [33] F. Farchioni, I. Hip and C.B. Lang, Phys. Lett. B443 (1998) 214.
- [34] F. Farchioni, I. Hip, C.B. Lang and M. Wohlgenannt, Nucl. Phys. B549 (1999) 364.
- [35] N. Cundy, M. Teper and U. Wenger, hep-lat/0203030.
- [36] G.P. Lepage and P. Mackenzie, Phys. Rev. D48 (1993) 2250.
- [37] C. Gattringer and I. Hip, Nucl. Phys. B541 (1999) 305.
- [38] M. Albanese *et al.*, Comput. Phys. Commun. 45 (1987) 345.
- [39] J. van den Eshof, A. Frommer, Th. Lippert, K. Schilling and H. van der Vorst, hep-lat/0202025; Nucl. Phys. B [Proc. Suppl.] 106 (2002) 1070.
- [40] H. Neuberger, Phys. Rev. Lett. 81 (1998) 4060.
- [41] R.G. Edwards, U.M. Heller and R. Narayanan, Nucl. Phys. B540 (1999) 457; Phys. Rev. D59 (1999) 094510. L. Giusti, C. Hoelbling and C. Rebbi, Phys. Rev. D64 (2001) 114508.
- [42] W.H. Press, S.A. Teukolsky, W.T. Vetterling and B.P. Flannery, “Numerical Recipes”, second edition, Cambridge University Press (Cambridge UK, 1992).
- [43] W. Bietenholz, I. Hip and K. Schilling, Nucl. Phys. B [Proc. Suppl.] 106 (2002) 829.
- [44] I. Horvath, Phys. Rev. Lett. 81 (1998) 4063; W. Bietenholz, hep-lat/9901005; I. Horvath, C. Balwe and R. Mendris, Nucl. Phys. B599 (2001) 283.
- [45] H. Neuberger, Phys. Rev. D61 (2000) 085015.
- [46] S. Chandrasekharan, Phys. Rev. D60 (1999) 074503. P.H. Damgaard, M.C. Diamantini, P. Hernández and K. Jansen, hep-lat/0112016.